



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MORTALITY—FOREIGN AND INSULAR—COUNTRIES AND CITIES
(Untabulated).

BRAZIL—Pernambuco.—Two weeks ended May 15, 1912. Population 225,000. Total number of deaths from all causes 408, including smallpox 34, tuberculosis 72, typhoid fever 2.

FRANCE—Nice.—Month of April, 1912. Population 168,185. Total number of deaths from all causes 239, including diphtheria 1, tuberculosis 38, typhoid fever 3. Cases reported: Diphtheria 1, scarlet fever 6, tuberculosis 3, typhoid fever 10.

Month of May, 1912. Total number of deaths from all causes 261, including diphtheria 1, tuberculosis 31, typhoid fever 2. Cases reported: Diphtheria 4, measles 4, scarlet fever 3, tuberculosis 1, typhoid fever 8.

GERMANY—Kehl.—Month of June, 1912. Population, 182,426. Total number of deaths from all causes 209, including diphtheria 2, measles 6, scarlet fever 2, tuberculosis 32, typhoid fever 1. Cases reported: Diphtheria, 34; scarlet fever, 44; tuberculosis, 8; typhoid fever, 3.

GREAT BRITAIN.—Week ended July 13, 1912.

England and Wales.—The deaths registered in 95 great towns correspond to an annual rate of 11.6 per 1,000 of the population, which is estimated at 17,639,816.

Ireland.—The deaths registered in 22 principal town districts correspond to an annual rate of 15.7 per 1,000 of the population, which is estimated at 1,157,014. The lowest rates were registered at Armagh, viz, 0.0, and Drogheda, viz, 0.0, and the highest rate was recorded at Kilkenny, viz, 34.7 per 1,000.

Scotland.—The deaths registered in 18 principal towns correspond to an annual rate of 13.4 per 1,000 of the population, which is estimated at 2,182,400. The lowest rates were registered at Govan, viz, 6.3, and Motherwell, viz, 6.3, per 1,000. The highest rate was recorded at Hamilton, viz, 23.8 per 1,000. The total number of deaths from all causes was 560, including diphtheria 4, measles 16, scarlet fever 3, typhoid fever 2.

SOUTH AFRICA—Johannesburg.—Two weeks ended June 24, 1912. Population, 237,220. Total number of deaths from all causes 156, including measles 2, tuberculosis 21, typhoid fever 2. Cases reported: Diphtheria, 15; scarlet fever, 12; typhoid fever, 13.

By authority of the Secretary of the Treasury:

RUPERT BLUE,
Surgeon General,
United States Public Health Service.

ADDITIONAL COPIES of this publication
may be procured from the SUPERINTEND-
ENT OF DOCUMENTS, Government Printing
Office, Washington, D. C., at 5 cents per copy

Subscription price, per volume - - - \$2
